

Insert the article title here

S. Cheng^{1,2} J.C. Ji¹ J. Zhou²

¹
²
 * E-mail:

ISSN 1751-8644
 doi: 0000000000
 www.ietdl.org

Abstract: The abstract text goes here.

1 Insert A head here

This demo file is intended to serve as a “starter file” for IMAIAI journal papers produced under L^AT_EX using imaiiai.cls v1.5e.

1.1 Insert B head here

Subsection text here.

1.1.1 Insert C head here: Subsubsection text here.

2 Equations

Sample equations.

$$\begin{aligned} \frac{\partial u(t, x)}{\partial t} &= Au(t, x) \left(1 - \frac{u(t, x)}{K} \right) - B \frac{u(t - \tau, x)w(t, x)}{1 + Eu(t - \tau, x)}, \\ \frac{\partial w(t, x)}{\partial t} &= \delta \frac{\partial^2 w(t, x)}{\partial x^2} - Cw(t, x) + D \frac{u(t - \tau, x)w(t, x)}{1 + Eu(t - \tau, x)}, \end{aligned} \tag{1}$$

$$\begin{aligned} \frac{dU}{dt} &= \alpha U(t)(\gamma - U(t)) - \frac{U(t - \tau)W(t)}{1 + U(t - \tau)}, \\ \frac{dW}{dt} &= -W(t) + \beta \frac{U(t - \tau)W(t)}{1 + U(t - \tau)}. \end{aligned} \tag{2}$$

$$\frac{\partial(F_1, F_2)}{\partial(c, \omega)} \Big|_{(c_0, \omega_0)} = \begin{vmatrix} \frac{\partial F_1}{\partial c} & \frac{\partial F_1}{\partial \omega} \\ \frac{\partial F_2}{\partial c} & \frac{\partial F_2}{\partial \omega} \end{vmatrix} \Big|_{(c_0, \omega_0)}$$

3 Enunciations

Theorem 1. Assume that $\alpha > 0, \gamma > 1, \beta > \frac{\gamma+1}{\gamma-1}$. Then there exists a small $\tau_1 > 0$, such that for $\tau \in [0, \tau_1)$, if c crosses $c(\tau)$ from the direction of to a small amplitude periodic traveling wave solution of (2.1), and the period of $(\check{u}^P(s), \check{w}^P(s))$ is

$$\check{T}(c) = c \cdot \left[\frac{2\pi}{\omega(\tau)} + O(c - c(\tau)) \right].$$

4 Figures & Tables

The output for figure is:

An example of a double column floating figure using two sub-figures. (The subfig.sty package must be loaded for this to work.) The subfigure \label commands are set within each subfloat

Fig. 1: Insert figure caption here

command, the \label for the overall figure must come after \caption. \hfil must be used as a separator to get equal spacing. The subfigure.sty package works much the same way, except \subfigure is used instead of \subfloat.

The output for table is:

Table 1 An Example of a Table

One	Two
Three	Four

5 Conclusion

The conclusion text goes here.

Acknowledgment

Insert the Acknowledgment text here.

6 References

- Vicsek T., Czirók A., Ben-Jacob E., Cohen I.: ‘Novel type of phase transition in a system of self-driven particles’, *Phys. Rev. Lett.*, 1995, **75**, pp. 1226–1229
- Jadbabaie A., Lin J., Morse A.S.: ‘Coordination of groups of mobile autonomous agents using nearest neighbor rules’, *IEEE Trans. Autom. Control*, 2003, **48**, pp. 988–1001
- Olfati-Saber R., Murray R.M.: ‘Consensus problems in networks of agents with switching topology and time-delays’, *IEEE Trans. Autom. Control*, 2004, **49**, pp. 1520–1533